

On Potentially $(K_5 - H)$ -graphic Sequences

Lili Hu and Chunhui Lai *

Department of Mathematics
Zhangzhou Teachers College, Zhangzhou
Fujian 363000, P. R. of China
jackey2591924@163.com(Lili Hu)
zjlaichu@public.zzptt.fj.cn(Chunhui Lai, Corresponding author)

Ping Wang †

Dept. of Math., Stats. and Computer Science
St. Francis Xavier University
Antigonish, NS, Canada, B2G 2W5
pwang@stfx.ca(Ping Wang)

Abstract

Let $K_m - H$ be the graph obtained from K_m by removing the edges set $E(H)$ of H where H is a subgraph of K_m . In this paper, we characterize the potentially $K_5 - P_4$ and $K_5 - Y_4$ -graphic sequences where Y_4 is a tree on 5 vertices and 3 leaves.

1 Introduction

We consider finite simple graphs. Any undefined notation follows that of Bondy and Murty [1]. An n -term non-increasing nonnegative integer sequence $\pi = (d_1, d_2, \dots, d_n)$ is said to be graphic if it is the degree sequence of a simple graph G of order n ; such a graph G is referred as a realization of π . Let $\sigma(\pi)$ the sum of all the terms of π , and let $[x]$ be the largest integer less than or equal to x . Let Y_4 denote a tree on 5 vertices and 3 leaves. A graphic sequence π is said to be potentially H -graphic if it has a realization G containing H as a subgraph. Let $G - H$ denote the graph obtained from G by removing the edges set $E(H)$ where H is a subgraph of G . In the degree sequence, r^t means

*Research is supported by NNSF of China(10271105) and by NSF of Fujian(Z0511034), Fujian Provincial Training Foundation for "Bai-Quan-Wan Talents Engineering", Project of Fujian Education Department and Project of Zhangzhou Teachers College.

†Research is support by NSERC

r repeats t times, that is, in the realization of the sequence there are t vertices of degree r .

In 1907, Mantel first proposed the problem of determining the maximum number of edges in a graph without containing 3-cycles. In general, this problem can be phrased as determining the maximum number of edges, denoted $ex(n, H)$, of a graph with n vertices not containing H as a subgraph. This area of research is called extremal graph theory. In terms of graphic sequences, the number $2ex(n, H) + 2$ is the minimum even integer l such that every n -term graphical sequence π with $\sigma(\pi) \geq l$ is forcibly H -graphical. In 1991, Erdős, Jacobson and Lehel [2] showed $\sigma(K_k, n) \geq (k-2)(2n-k+1)+2$ and conjectured that the equality holds. In the same paper, they proved the conjecture is true for the case $k=3$ and $n \geq 6$. The cases $k=4$ and 5 were proved separately in [3], [11] and [12]. Based on linear algebraic techniques, Li, Song and Luo [13] proved the conjecture true for $k \geq 6$ and $n \geq \binom{k}{2} + 3$. Recently, Ferrara, Gould and Schmitt proved the conjecture [4] and they also determined in [5] $\sigma(F_k, n)$ where F_k denotes the graph of k triangles intersecting at exactly one common vertex.

In 1999, Gould, Jacobson and Lehel [3] considered the following generalized problem: determine the smallest even integer $\sigma(H, n)$ such that every n -term positive graphic sequence $\pi = (d_1, d_2, \dots, d_n)$ with $\sigma(\pi) \geq \sigma(H, n)$ has a realization G containing H as a subgraph. They proved $\sigma(pK_2, n) = (p-1)(2n-p)+2$ for $p \geq 2$ and $\sigma(C_4, n) = 2\lceil \frac{3n-1}{2} \rceil$ for $n \geq 4$. Lai [8] determined $\sigma(K_4 - e, n)$ for $n \geq 4$. Yin, Li, and Mao [19] determined $\sigma(K_{r+1} - e, n)$ for $r \geq 3$ and $r+1 \leq n \leq 2r$ and $\sigma(K_5 - e, n)$ for $n \geq 5$, and Yin and Li [18] further determined $\sigma(K_{r+1} - e, n)$ for $r \geq 2$ and $n \geq 3r^2 - r - 1$. Moreover, Yin and Li in [18] also gave two sufficient conditions for a sequence $\pi \in GS_n$ to be potentially $(K_{r+1} - e)$ -graphic. Yin [21] determined $\sigma(K_{r+1} - K_3, n)$ for $r \geq 3$ and $n \geq 3r+5$. Lai [9] determined $\sigma(K_5 - P_3, n)$ and $\sigma(K_5 - P_4, n)$ for $n \geq 5$. Lai and Hu [10] determined $\sigma(K_{r+1} - H, n)$ for $n \geq 4r+10$, $r \geq 3$, $r+1 \leq k \leq 4$ and H be a graph on k vertices which containing a tree on 4 vertices but not containing a cycle on 3 vertices and $\sigma(K_{r+1} - P_2, n)$ for $n \geq 4r+8$, $r \geq 3$.

A harder question is to characterize the potentially H -graphic sequences without zero terms. That is, finding necessary and sufficient conditions for a sequence to be a H -graphic sequence. Luo [15] characterized the potentially C_k -graphic sequences for each $k=3, 4$ and 5 . Recently, in [16], Luo and Warner also characterized the potentially K_4 -graphic sequences. Eschen and Niu [17] characterized the potentially $(K_4 - e)$ -graphic sequences. Hu and Lai [6] characterized the potentially $(K_5 - C_4)$ -graphic sequences. Yin and Chen [20] characterized the potentially $K_{r,s}$ -graphic sequences for $r=2, s=3$ and $r=2, s=4$, where $K_{r,s}$ is an $r \times s$ complete bipartite graph.

In attempt to completely characterize the potentially $K_5 - H$ - graphic sequences, we will characterize the potentially $K_5 - P_4$ and $K_5 - Y_4$ - graphic sequences in this paper.

Let $\pi = (d_1, d_2, \dots, d_n)$ be a nonincreasing positive integer sequence. We write $m(\pi)$ and $h(\pi)$ to denote the largest positive terms of π and the smallest positive terms of π , respectively. $\pi'' = (d_1-1, d_2-1, \dots, d_{d_n}-1, d_{d_n+1}, \dots, d_{n-1})$

is the residual sequence obtained by laying off d_n from π . We denote $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$ where $d'_1 \geq d'_2 \geq \dots \geq d'_{n-1}$ is a rearrangement of the $n-1$ terms in π' . We denote π'' the residual sequence obtained by laying off d_n from π and all the graphic sequences have no zero terms. We need the following results.

Theorem 1.1. [3] *If $\pi = (d_1, d_2, \dots, d_n)$ is a graphic sequence with a realization G containing H as a subgraph, then there exists a realization G' of π containing H as a subgraph so that the vertices of H have the largest degrees of π .*

Theorem 1.2. [14] *If $\pi = (d_1, d_2, \dots, d_n)$ is a sequence of nonnegative integers with $1 \leq m(\pi) \leq 2$, $h(\pi) = 1$ and even $\sigma(\pi)$, then π is graphic.*

Theorem 1.3. [7] *π is graphic if and only if π' is graphic.*

The following corollary is obvious.

Corollary 1.4. *Let H be a simple graph. If π' is potentially H -graphic, then π is potentially H -graphic.*

2 Main Theorems

Theorem 2.1. *Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $(K_5 - P_4)$ -graphic if and only if the following conditions hold:*

1. $d_2 \geq 3$.
2. $d_5 \geq 2$.
3. $\pi \neq (n-1, k, 2^t, 1^{n-2-t})$ where $n \geq 5$, $k, t = 3, 4, \dots, n-2$, and, k and t have different parities.
4. For $n \geq 5$, $\pi \neq (n-k, k+i, 2^i, 1^{n-i-2})$ where $i = 3, 4, \dots, n-2k$ and $k = 1, 2, \dots, \lfloor \frac{n-1}{2} \rfloor - 1$.
5. If $n = 6, 7$, then $\pi \neq (3^2, 2^{n-2})$.

Proof. First we show the conditions (1)-(5) are necessary conditions for π to be potentially $(K_5 - P_4)$ -graphic. Assume that π is potentially $(K_5 - P_4)$ -graphic. (1), (2) and (5) are obvious. If $\pi = (n-1, k, 2^t, 1^{n-2-t})$ is potentially $(K_5 - P_4)$ -graphic, then according to Theorem 1.1, there exists a realization G of π containing $K_5 - P_4$ as a subgraph so that the vertices of $K_5 - P_4$ have the largest degrees of π . Therefore, the sequence $\pi^* = (n-4, k-3, 2^{t-3}, 1^{n-2-t})$ obtained from $G - (K_5 - P_4)$ must be graphic. Since the edge between two vertices with degree $n-4$ and $k-3$ has been removed from the realization of π^* , thus, $\Delta(G - (K_5 - P_4)) \leq n-5$, a contradiction. Hence, (3) holds. If $\pi = (n-k, k+i, 2^i, 1^{n-i-2})$ is potentially $(K_5 - P_4)$ -graphic, then according to Theorem 1.1, there exists a realization G of π containing $K_5 - P_4$ as a subgraph so that the vertices of $K_5 - P_4$ have the largest degrees of π . Therefore, the

sequence $\pi^* = (n - k - 3, k + i - 3, 2^{i-3}, 1^{n-i-2})$ obtained from $G - (K_5 - P_4)$ must be graphic and there is no edge between two vertices with degree $n - k - 3$ and $k + i - 3$ in the realization of π^* . Let G^* be a realization of π^* , and, $d_{G^*}(x) = n - k - 3$ and $d_{G^*}(y) = k + i - 3$. Consider a partition of G^* where $X = \{x, y\}$ and $Y = V(G^*) - \{x, y\}$. It follows that the number of edges between X and Y equals $(n - k - 3) + (k + i - 3) \leq 2(i - 3) + (n - i - 2)$, that is, $[(n - k - 3) + (k + i - 3)] - [2(i - 3) + (n - i - 2)] = 2 \leq 0$, a contradiction. Hence, (4) holds.

Now we turn to show the conditions (1)-(5) are sufficient conditions for π to be potentially $(K_5 - P_4)$ -graphic. Suppose the graphic sequence π satisfies the conditions (1) to (5). Our proof is by induction on n . We first prove the base case where $n = 5$. Since $\pi \neq (4^2, 2^3)$, then π must be one of the following sequences: (4^5) , $(4^3, 3^2)$, $(4^2, 3^2, 2)$, $(4, 3^4)$, $(4, 3^2, 2^2)$, $(3^4, 2)$, $(3^2, 2^3)$. It is easy to check that all of these are potentially $(K_5 - P_4)$ -graphic. Now we assume that the sufficiency holds for $n - 1$ ($n \geq 6$). We will prove π is potentially $(K_5 - P_4)$ -graphic.

Case 1: $\pi' = (3^2, 2^4)$

Clearly, $n = 7$ and π must be one of the following sequences $(4^2, 2^5)$, $(4, 3^2, 2^4)$, $(3^4, 2^3)$, $(4, 3, 2^4, 1)$ or $(3^3, 2^3, 1)$. It is easy to check that all of these are potentially $(K_5 - P_4)$ -graphic.

Case 2: $\pi' = (3^2, 2^5)$

Clearly, $n = 8$ and π must be one of the following sequences $(4^2, 2^6)$, $(4, 3^2, 2^5)$, $(3^4, 2^4)$, $(4, 3, 2^5, 1)$ or $(3^3, 2^4, 1)$. It is easy to check that all of these are potentially $(K_5 - P_4)$ -graphic.

Case 3: $d_n \geq 3$

Clearly, π' satisfies the assumption, and thus, by the induction hypothesis, π' is potentially $(K_5 - P_4)$ -graphic, and hence so is π . In the following, we only consider the cases $d_n = 1$ or $d_n = 2$.

Case 4: $\pi' = (n - 2, k, 2^t, 1^{n-3-t})$ where $n - 1 \geq 5$, $k, t = 3, 4, \dots, n - 3$, and, k and t have different parities

If $d_n = 2$, then $\pi' = (n - 2, k, 2^{n-3})$. If $k \geq 4$, then $\pi = (n - 1, k + 1, 2^{n-2})$ which contradicts condition (3). If $k = 3$, that is $\pi' = (n - 2, 3, 2^{n-3})$, then $\pi = (n - 1, 4, 2^{n-2})$ or $\pi = (n - 1, 3^2, 2^{n-3})$. But $\pi = (n - 1, 4, 2^{n-2})$ contradicts condition (3), thus $\pi = (n - 1, 3^2, 2^{n-3})$ where n is odd. We will show $\pi = (n - 1, 3^2, 2^{n-3})$ is potentially $(K_5 - P_4)$ -graphic. In other words, we would like to show $\pi_1 = (n - 4, 2^{n-5}, 1)$ is graphic. It suffices to show $\pi_2 = (1^{n-5})$ where $n \geq 7$ is graphic. By $\sigma(\pi_2)$ being even and Theorem 1.2, π_2 is graphic. Thus, $\pi = (n - 1, 3^2, 2^{n-3})$ is potentially $(K_5 - P_4)$ -graphic.

If $d_n = 1$, then $\pi = (n - 1, k, 2^t, 1^{n-2-t})$ which contradicts condition (3).

Case 5: $\pi' = (n - 1 - k, k + i, 2^i, 1^{n-i-3})$ where $i = 3, 4, \dots, n - 1 - 2k$ and $k = 1, 2, \dots, \lceil \frac{n}{2} \rceil - 2$.

If $d_n = 2$, then $n - i - 3 = 0$ and $\pi = (n - k, k + i + 1, 2^{i+1})$ and it contradicts condition (4).

If $d_n = 1$ and $n - 1 - k = k + i + 1$, then $\pi = (n - k, k + i, 2^i, 1^{n-i-2})$ or $\pi = ((n - 1 - k)^2, 2^i, 1^{n-i-2})$, both of these are contradict to condition (4). If $d_n = 1$ and $n - 1 - k = k + i$ or $n - 1 - k \geq k + i + 2$, then $\pi = (n - k, k + i, 2^i, 1^{n-i-2})$ which also contradicts condition (4).

Case 6: $d_n = 2$, $\pi' \neq (n - 2, k, 2^{n-3})$, $\pi' \neq (n - 1 - k, n + k - 3, 2^{n-3})$, $\pi' \neq (3^2, 2^4)$, and $\pi' \neq (3^2, 2^5)$

Consider $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$. Since $d_2 \geq 3$, we have $d'_{n-1} \geq 2$. If $d'_2 \geq 3$, then π' satisfies the assumption. Thus, π' is potentially $(K_5 - P_4)$ -graphic. Hence, we may assume $d'_2 = 2$, that is, $d_2 = 3$ and $d_3 = d_4 = \dots = d_n = 2$. It follows $\pi = (d_1, 3, 2^{n-2})$. Since $\sigma(\pi)$ is even, d_1 must be odd. If $d_1 = 3$, then $\pi = (3^2, 2^{n-2})$. Since $\pi \neq (3^2, 2^4)$ and $\pi \neq (3^2, 2^5)$, we have $n \geq 8$. We will show π is potentially $(K_5 - P_4)$ -graphic. It suffices to show $\pi_1 = (2^{n-5})$ is graphic. Clearly, C_{n-5} is a realization of π_1 . If $d_1 \geq 5$, since $\pi \neq (n - 1, 3, 2^{n-2})$, we have $d_1 \leq n - 2$. We will prove $\pi = (d_1, 3, 2^{n-2})$ where $d_1 \geq 5$ and $n \geq d_1 + 2$ is potentially $(K_5 - P_4)$ -graphic. We would like to show $\pi_1 = (d_1 - 3, 2^{n-5})$ is graphic. It suffices to show $\pi_2 = (2^{n-d_1-2}, 1^{d_1-3})$ is graphic. Since $\sigma(\pi_2)$ is even, π_2 is graphic by Theorem 1.2. Thus, $\pi = (d_1, 3, 2^{n-2})$ is potentially $(K_5 - P_4)$ -graphic.

Case 7: $d_n = 1$, $\pi' \neq (n - 2, k, 2^t, 1^{n-3-t})$, $\pi' \neq (n - 1 - k, k + i, 2^i, 1^{n-i-3})$, $\pi' \neq (3^2, 2^4)$, and $\pi' \neq (3^2, 2^5)$

Consider $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$. Since $d_2 \geq 3$ and $d_5 \geq 2$, we have $d'_1 \geq 3$ and $d'_5 \geq 2$. If $d'_2 \geq 3$, then π' satisfies the assumption. Thus, π' is potentially $(K_5 - P_4)$ -graphic. Hence, we may assume $d'_2 = 2$, that is, $d_1 = d_2 = 3$ and $d_3 = d_4 = d_5 = 2$. Thus $\pi = (3^2, 2^t, 1^{n-2-t})$ where $t \geq 3$ and $n - 2 - t \geq 1$. Since $\sigma(\pi)$ is even, $n - 2 - t$ must be even. We will prove π is potentially $(K_5 - P_4)$ -graphic. It suffices to show $\pi_1 = (2^{t-3}, 1^{n-2-t})$ is graphic. Since $\sigma(\pi_1)$ is even, π_1 is graphic by Theorem 1.2 and, in turn, π is potentially $(K_5 - P_4)$ -graphic.

This completes the proof. $\square$

Theorem 2.2. *Let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence with $n \geq 5$. Then π is potentially $(K_5 - Y_4)$ -graphic if and only if the following conditions hold:*

1. $d_3 \geq 3$.
2. $d_4 \geq 2$.
3. $\pi \neq (3^6)$.

Proof. Assume that π is potentially $(K_5 - Y_4)$ -graphic. In this case the necessary conditions (1) to (3) are obvious.

Now we prove the sufficient conditions. Suppose the graphic sequence π satisfies the conditions (1) to (3). Our proof is by induction on n . We first prove the base case where $n = 5$. In this case, π is one of the following sequences:

(4^5) , $(4^3, 3^2)$, $(4^2, 3^2, 2)$, $(4, 3^4)$, $(4, 3^3, 1)$, $(4, 3^2, 2^2)$, $(3^4, 2)$, or $(3^3, 2, 1)$. It is easy to check that all of these are potentially $(K_5 - Y_4)$ -graphic. Now suppose the sufficiency holds for $n - 1$ ($n \geq 6$), and let $\pi = (d_1, d_2, \dots, d_n)$ be a graphic sequence which satisfies (1) to (3). We will prove π is potentially $(K_5 - Y_4)$ -graphic.

Case 1: $\pi' = (3^6)$

We have $n = 7$ and π is one of the following sequences $(4^3, 3^4)$, $(4^2, 3^4, 2)$ or $(4, 3^5, 1)$. It is easy to check that all of these are potentially $(K_5 - Y_4)$ -graphic.

Case 2: $d_n \geq 3$ and $\pi' \neq (3^6)$

Clearly, $d'_4 \geq 2$. If $d_3 \geq 4$, then $d'_3 \geq 3$. If $d_3 = \dots = d_n = 3$ and $n \geq 6$, $d'_3 \geq 3$. It follows conditions (1) and (2) hold. Thus, by the induction hypothesis, π' is potentially $(K_5 - Y_4)$ -graphic. Therefore, π is potentially $(K_5 - Y_4)$ -graphic by Corollary 1.4. In the following, we only consider the cases where $d_n = 2$ or $d_n = 1$.

Case 3: $d_n = 2$ and $\pi' \neq (3^6)$

Consider $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$. Since $d_3 \geq 3$ and $d_n = 2$, we have $d'_1 \geq 3$ and $d'_{n-1} \geq 2$. If $d'_3 \geq 3$, then π' satisfies the assumption and it follows π' is potentially $(K_5 - Y_4)$ -graphic. Therefore, π is potentially $(K_5 - Y_4)$ -graphic by Corollary 1.4. Hence, we may assume $d'_3 = 2$. We will proceed with the following two cases $d_1 \geq 4$ and $d_1 = 3$.

Subcase 1: $d_1 \geq 4$

It suffices to consider the case where $d_2 = d_3 = 3$ and $d_4 = d_5 = \dots = d_n = 2$. That is, $\pi = (d_1, 3^2, 2^{n-3})$. Since $\sigma(\pi)$ is even, d_1 must be even. We will prove π is potentially $(K_5 - Y_4)$ -graphic. It is enough to show $\pi_1 = (d_1 - 3, 2^{n-5}, 1)$ is graphic. If $d_1 = n - 1$, then $\pi_1 = (n - 4, 2^{n-5}, 1)$. It suffices to show $\pi_2 = (1^{n-5})$ is graphic. Since $\sigma(\pi_2)$ is even, π_2 is graphic by Theorem 1.2. If $d_1 \leq n - 2$, it suffices to show $\pi_2 = (2^{n-2-d_1}, 1^{d_1-2})$ (or $\pi_2 = (2^{n-1-d_1}, 1^{d_1-4})$) is graphic. Similarly, one can show π_2 is graphic. Thus, $\pi_1 = (d_1 - 3, 2^{n-5}, 1)$ is graphic and, in turn, π is potentially $(K_5 - Y_4)$ -graphic.

Subcase 2: $d_1 = 3$

It suffices to consider the case where $d_1 = d_2 = d_3 = d_4 = 3$ and $d_5 = \dots = d_n = 2$. That is, $\pi = (3^4, 2^{n-4})$. We will prove π is potentially $(K_5 - Y_4)$ -graphic. It is enough to show $\pi_1 = (2^{n-5}, 1^2)$ is graphic. Since $\sigma(\pi_1)$ is even, π_1 is graphic by Theorem 1.2 and, in turn, π is potentially $(K_5 - Y_4)$ -graphic.

Case 4: $d_n = 1$ and $\pi' \neq (3^6)$

Consider $\pi' = (d'_1, d'_2, \dots, d'_{n-1})$. Since $d_3 \geq 3$ and $d_4 \geq 2$, we have $d'_2 \geq 3$ and $d'_4 \geq 2$. If $d'_3 \geq 3$, then π' satisfies the assumptions and it follows π' is potentially $(K_5 - Y_4)$ -graphic. Therefore, π is potentially $(K_5 - Y_4)$ -graphic by Corollary 1.4. Hence, we may assume $d'_3 = 2$. It suffices to consider the case where $d_1 = d_2 = d_3 = 3$ and $d_4 = 2$. That is, $\pi = (3^3, 2^t, 1^{n-3-t})$ where $t \geq 1$

and $n - 3 - t \geq 1$. Since $\sigma(\pi)$ is even, $n - t$ must be even. We will prove π is potentially $(K_5 - Y_4)$ -graphic. It is enough to show $\pi_1 = (2^{t-2}, 1^{n-2-t})$ is graphic when $t \geq 2$. Since $\sigma(\pi_1)$ is even, π_1 is graphic by Theorem 1.2. If $t = 1$, then $\pi = (3^3, 2, 1^{n-4})$. Similarly we can show $\pi_2 = (1^{n-5})$ is graphic and, in turn, π potentially $(K_5 - Y_4)$ -graphic.

This completes the proof. $\square$

In the remaining of this section, we will use the above two theorems to find exact values of $\sigma(K_5 - P_4, n)$ and $\sigma(K_5 - Y_4, n)$. Note that the value of $\sigma(K_5 - P_4, n)$ was determined by Lai in [9] so a much simpler proof is given here.

Corollary 2.3. ([9]) For $n \geq 5$, $\sigma(K_5 - P_4, n) = 4n - 4$.

Proof. First we claim $\sigma(K_5 - P_4, n) \geq 4n - 4$ for $n \geq 5$. We would like to show there exists π_1 with $\sigma(\pi_1) = 4n - 6$ such that π_1 is not potentially $(K_5 - P_4)$ -graphic. Let $\pi_1 = ((n-1)^2, 2^{n-2})$. It is easy to see that $\sigma(\pi_1) = 4n - 6$ and π_1 is not potentially $(K_5 - P_4)$ -graphic by Theorem 2.1.

Now we show if π is an n -term ($n \geq 5$) graphical sequence with $\sigma(\pi) \geq 4n - 4$, then there exists a realization of π containing a $K_5 - P_4$. If $d_5 = 1$, then $\sigma(\pi) = d_1 + d_2 + d_3 + d_4 + (n - 4)$. Let X be the four vertices of the largest degrees of G and $Y = V(G) - X$. Since there are at most six edges in X , $d_1 + d_2 + d_3 + d_4 \leq 12 + |E(X, Y)| \leq 12 + (n - 4) = n + 8$. This leads to $\sigma(\pi) \leq 2n + 4 < 4n - 4$, a contradiction. Thus, $d_5 \geq 2$. If $d_2 \leq 2$, then $\sigma(\pi) \leq d_1 + 2(n - 1) \leq 3n - 3 < 4n - 4$, a contradiction. Thus, $d_2 \geq 3$. Since $\sigma(\pi) \geq 4n - 4$, then π is not one of the following: $(3^2, 2^4)$, $(3^2, 2^5)$, and $(n-1, k, 2^t, 1^{n-2-t})$ where $n \geq 6$ and $k, t = 3, 4, \dots, n-2$, $(n-k, k+i, 2^i, 1^{n-i-2})$ where $i = 3, 4, \dots, n-2k$ and $k = 1, 2, \dots, \lfloor \frac{n-1}{2} \rfloor - 1$. Thus, π satisfies the conditions (1) to (5) in Theorem 2.1. Therefore, π is potentially $(K_5 - P_4)$ -graphic by Theorem 2.1. $\square$

Corollary 2.4. For $n \geq 5$, $\sigma(K_5 - Y_4, n) = 4n - 4$.

Proof. First we claim $\sigma(K_5 - Y_4, n) \geq 4n - 4$ if $n \geq 5$. We would like to show there exists π_1 with $\sigma(\pi_1) = 4n - 6$, such that π_1 is not potentially $(K_5 - Y_4)$ -graphic. Let $\pi_1 = ((n-1)^2, 2^{n-2})$. It is easy to see that $\sigma(\pi_1) = 4n - 6$ and π_1 is not potentially $(K_5 - Y_4)$ -graphic by Theorem 2.2.

Now we show if π is an n -term ($n \geq 5$) graphical sequence with $\sigma(\pi) \geq 4n - 4$, then there exists a realization of π containing a $K_5 - Y_4$. If $d_4 = 1$, then $\sigma(\pi) = d_1 + d_2 + d_3 + (n - 3)$. By the similar argument used in the above corollary, we have $d_1 + d_2 + d_3 \leq 6 + (n - 3) = n + 3$. This leads to $\sigma(\pi) \leq 2n < 4n - 4$, a contradiction. Thus, $d_4 \geq 2$. Similarly, if $d_3 \leq 2$, then $\sigma(\pi) \leq d_1 + d_2 + 2(n - 2) \leq 2(n - 1) + 2(n - 2) = 4n - 6 < 4n - 4$, a contradiction. Thus, $d_3 \geq 3$. Since $\sigma(\pi) \geq 4n - 4$, then $\pi \neq (3^6)$. Thus, π satisfies the conditions (1) to (3) in Theorem 2.2. Therefore, π is potentially $(K_5 - Y_4)$ -graphic by Theorem 2.2. $\square$

References

- [1] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications, The Macmillan Press Ltd., 1976.
- [2] P.Erdős, M.S. Jacobson and J. Lehel, Graphs realizing the same degree sequences and their respective clique numbers, in Graph Theory, Combinatorics and Application, Vol. 1 (Y. Alavi et al., eds.), John Wiley and Sons, Inc., New York, (1991): 439-449.
- [3] R.J. Gould, M.S. Jacobson and J. Lehel, Potentially G -graphic degree sequences, Combinatorics, Graph Theory and Algorithms, Vol. 2 (Y. Alavi et al., eds.), New Issues Press, Kalamazoo, MI, (1999): 451-460.
- [4] Ferrara, M., Gould, R., and Schmitt, J., Potentially K_s^t -graphic degree sequences, submitted.
- [5] Ferrara, M., Gould, R., and Schmitt, J., Graphic sequences with a realization containing a friendship graph, accepted by Ars Combinatoria.
- [6] Lili Hu and Chunhui Lai, On potentially $K_5 - C_4$ -graphic sequences, accepted by Ars Combinatoria. math.CO/0701023
- [7] D.J. Kleitman and D.L. Wang, Algorithm for constructing graphs and digraphs with given valences and factors, Discrete Math., 6(1973): 79-88.
- [8] Chunhui Lai, A note on potentially $K_4 - e$ graphical sequences, Australasian Journal of Combinatorics 24(2001), 123-127. math.CO/0308105
- [9] Chunhui Lai, An extremal problem on potentially $K_m - P_k$ -graphic sequences, accepted by International Journal of Pure and Applied Mathematics. math.CO/0409466
- [10] Chunhui Lai and Lili Hu, An extremal problem on potentially $K_{r+1} - H$ -graphic sequences, accepted by Ars Combinatoria. math.CO/0603265
- [11] Jiong-Sheng Li and Zi-Xia Song, An extremal problem on the potentially P_k -graphic sequences, The International Symposium on Combinatorics and Applications, June 28-30, 1996 (W.Y.C. Chen et. al., eds.) Tianjin, Nankai University (1996): 269-276.
- [12] Jiong-Sheng Li and Zi-Xia Song, The smallest degree sum that yields potentially P_k -graphical sequences, J. of Graph Theory, 29(1998): 63-72.
- [13] Jiong-sheng Li, Zi-Xia Song and Rong Luo, The Erdős-Jacobson-Lehel conjecture on potentially P_k -graphic sequence is true, Science in China(Series A), 41(5)(1998): 510-520.
- [14] Jiong-sheng Li and Jianhua Yin, A variation of an extremal theorem due to Woodall, Southeast Asian Bulletin of Math., 25(2001): 427-434.

- [15] Rong Luo, On potentially C_k -graphic sequences, *Ars Combinatoria* 64(2002): 301-318.
- [16] Rong Luo and Morgan Warner, On potentially K_k -graphic sequences, *Ars Combinatoria* 75(2005): 233-239.
- [17] Elaine M. Eschen and Jianbing Niu, On potentially $K_4 - e$ -graphic sequences, *Australasian Journal of Combinatorics*, 29(2004): 59-65.
- [18] Jianhua Yin and Jiongsheng Li, Two sufficient conditions for a graphic sequence to have a realization with prescribed clique size, *Discrete Math.*, 301(2005): 218-227.
- [19] Jianhua Yin, Jiongsheng Li and Rui Mao, An extremal problem on the potentially $K_{r+1} - e$ -graphic sequences, *Ars Combinatoria* 74(2005), 151-159.
- [20] Jianhua Yin and Gang Chen, On potentially $K_{r_1, r_2, \dots, r_m}$ -graphic sequences, preprint.
- [21] Mengxiao Yin, The smallest degree sum that yields potentially $K_{r+1} - K_3$ -graphic sequences, accepted by *Acta Mathematicae Applicatae Sinica*.